Noname manuscript No.
(will be inserted by the editor)

arXiv:astro-ph/0608372v1 17 Aug 2006

A. M. Beloborodov · C. Thompson

Magnetar Corona

Received: date / Accepted: date

Abstract Persistent high-energy emission of magnetars is produced by a plasma corona around the neutron star, with total energy output of $\sim 10^{36}$ erg/s. The corona forms as a result of occasional starquakes that twist the external magnetic field of the star and induce electric currents in the closed magnetosphere. Once twisted, the magnetosphere cannot untwist immediately because of its self-induction. The self-induction electric field lifts particles from the stellar surface, accelerates them, and initiates avalanches of pair creation in the magnetosphere. The created plasma corona maintains the electric current demanded by curl $\mathbf{B}$ and regulates the self-induction e.m.f. by screening. This corona persists in dynamic equilibrium: it is continually lost to the stellar surface on the light-crossing time $\sim 10^{-4}$ s and replenished with new particles. In essence, the twisted magnetosphere acts as an accelerator that converts the toroidal field energy to particle kinetic energy. The voltage along the magnetic field lines is maintained near threshold for ignition of pair production, in the regime of self-organized criticality. The voltage is found to be about ~ 1 GeV which is in agreement with the observed dissipation rate $\sim 10^{36}$ erg/s. The coronal particles impact the solid crust, knock out protons, and regulate the column density of the hydrostatic atmosphere of the star. The transition layer between the atmosphere and the corona is the likely source of the observed 100 keV emission from magnetars. The corona also emits curvature radiation up to 10^{14} Hz and can supply the observed IR-optical luminosity.

Keywords plasmas · stars: coronae, magnetic fields, neutron · X-rays: stars

A. M. Beloborodov
Physics Department and Columbia Astrophysics Laboratory, Columbia University, 538 W 120th Street, New York, US
E-mail: amb@phys.columbia.edu

C. Thompson
Canadian Institute for Theoretical Astrophysics, University of Toronto, 60 St. George Street, Toronto, Canada; E-mail: thompson@cita.utoronto.ca

1 Introduction

At least 10% of all neutron stars are born as magnetars, with magnetic fields $B > 10^{14}$ G. Their activity is powered by the decay of the ultrastrong field and lasts about 10^4 years. They are observed at this active stage as either Soft Gamma Repeaters (SGRs) or Anomalous X-ray Pulsars (AXPs) [1].

Besides the sporadic X-ray outbursts, a second, persistent, form of activity has been discovered by studying the emission spectra of magnetars. Until recently, the spectrum was known to have a thermal component with temperature $k_{\rm B}T \sim 0.5$ keV, interpreted as blackbody emission from the star's surface. The soft X-ray spectrum also showed a tail at $2-10$ keV with photon index $\Gamma = 2-4$. This deviation from pure surface emission already suggested that energy is partially released above the star's surface. Recently, observations by RXTE and INTEGRAL have revealed even more intriguing feature: magnetars are bright, persistent sources of 100 keV X-rays [2]. This high-energy emission forms a separate component of the magnetar spectrum. It becomes dominant above 10 keV, has a very hard spectrum, $\Gamma \simeq 1$, and peaks above 100 keV where the spectrum is unknown. Its luminosity, $L \sim 10^{36}$ erg s^{-1}, even exceeds the thermal luminosity from the star's surface. The observed hard X-rays can be emitted only in the exterior of the star and demonstrate the presence of an active plasma corona.

This corona can form as a result of starquakes that shear the magnetar crust and its external magnetic field. The persistence of magnetospheric twists can explain several observed properties of magnetars. Thompson, Lyutikov, & Kulkarni [3] investigated the observational consequences using a static, force-free model, idealizing the magnetosphere as a globally twisted dipole. They showed that a twist affects the spindown rate of the neutron star: it causes the magnetosphere to flare out slightly from a pure dipole configuration, thereby increasing the braking torque acting on the star. Although the calculations of force-free configurations are independent of the plasma behavior in the magnetosphere, they rely on the presence of plasma that can conduct the required current $\mathbf{j}_B = (c/4\pi)\nabla \times \mathbf{B}$. This requires a minimum particle density $n_c = j_B/ec$. Thompson et al. [3] showed that even this minimum density can modify the stellar radiation by multiple resonant scattering.

The problem of plasma dynamics in the closed twisted magnetosphere has been formulated in [4], and a simple solution has been found to this problem. It allows one to understand the observed energy output of the magnetar corona and the evolution of magnetic twists. It is worth mentioning that plasma behavior around neutron stars has been studied for decades in the context of radio pulsars, and that problem remains unsolved. The principle difference with radio pulsars is that their activity is caused by *rotation*, and dissipation occurs on *open magnetic lines* that connect the star to its light cylinder. Electron-positron pairs are then created on open field lines [5,6]. By contrast, the formation of a corona around a magnetar does not depend on its rotation. All observed magnetars are slow rotators ($\Omega = 2\pi/P \sim 1$ Hz), and their rotational energy is unable to feed the observed coronal emission. Practically the entire plasma corona is immersed in the *closed* magnetosphere and its heating must be caused by some form of dissipation on the closed field lines. It is this closure that facilitates the corona problem by providing both boundary conditions at the two footpoints of a field line.

The basic questions that one would like to answer are as follows. How is the magnetosphere populated with plasma? The neutron star surface has a temperature $k_{\rm B}T \lesssim 1$ keV and the scale-height of its atmospheric layer (if it exists) is only a few cm. How is the plasma supplied above this layer and what type of particles populate the corona? How is the corona heated, and what are the typical energies of the particles? If the corona conducts a current associated with a magnetic twist $\nabla \times \mathbf{B} \neq 0$, how rapid is the dissipation of this current, i.e. what is the lifetime of the twist? Does its decay imply the disappearance of the corona?

An outline of the proposed model is as follows. The key agent in corona formation is an electric field $E_\parallel$ parallel to the magnetic field. $E_\parallel$ is generated by the self-induction of the gradually decaying current and in essence measures the rate of the decay. It determines the heating rate of the corona via Joule dissipation. If $E_\parallel = 0$ then the corona is not heated and, being in contact with the cool stellar surface, it will have to condense to a thin surface layer with $k_{\rm B}T \lesssim 1$ keV. The current-carrying particles cannot flow upward against gravity unless a force $eE_\parallel$ drives it. On the other hand, when $E_\parallel$ exceeds a critical value, $e^\pm$ avalanches are triggered in the magnetosphere, and the created pairs screen the electric field. This leads to a "bottleneck" for the decay of a magnetic twist, which implies a slow decay.

Maintenance of the corona and the slow decay of the magnetic twist are intimately related because both are governed by $E_\parallel$. In order to find $E_\parallel$, one can use Gauss' law $\nabla \cdot \mathbf{E} = 4\pi\rho$ where ρ is the net charge density of the coronal plasma. This constraint implies that $\mathbf{E}$ and ρ must be found self-consistently. The problem turns out to be similar to the classical double-layer problem of plasma physics with a new essential ingredient: $e^\pm$ creation.

A direct numerical experiment can be designed that simulates the creation and behavior of the plasma in the magnetosphere. The experiment shows how the plasma and electric field self-organize to maintain the time-average magnetospheric current $\bar{\mathbf{j}} = \mathbf{j}_B$ demanded by the magnetic field, $(4\pi/c)\bar{\mathbf{j}} = \nabla \times \mathbf{B}$. The electric current admits no steady state on short timescales and keeps fluctuating, producing $e^\pm$ avalanches. This state may be described as a self-organized criticality. Pair creation is found to provide a robust mechanism for limiting the voltage along the magnetic lines to $e\Phi_e \lesssim 1$ GeV and regulate the observed luminosity of the corona.

2 Mechanism of corona formation

A tightly wound-up magnetic field is assumed to exist inside magnetars [7]. The internal toroidal field can be much stronger than the external large-scale dipole component. The essence of magnetar activity is the transfer of magnetic helicity from the interior of the star to its exterior, where it dissipates. This involves rotational motions of the crust, which inevitably twist the external magnetosphere anchored to the stellar surface. The magnetosphere is probably twisted in a catastrophic way, when the internal magnetic field breaks the crust and rotates it. Such starquakes are associated with observed X-ray bursts [8]. The most interesting effect of a starquake for us here is the partial release of the winding of the internal magnetic field into the exterior, i.e., the injection of magnetic helicity into the magnetosphere.

Since the magnetic field is energetically dominant in the magnetosphere, it must relax there to a force-free configuration with $\mathbf{j} \times \mathbf{B} = 0$. Electric current $\mathbf{j}$ initiated by a starquake flows along the magnetic field lines to the exterior of the star, reaches the top of the field line, and comes back to the star at the other footpoint, then enters the atmosphere and the crust, following the magnetic field lines ($\mathbf{j} \parallel \mathbf{B}$). Sufficiently deep in the crust the current flows across the field lines and changes direction so that it can flow back to the twisted footpoint through the magnetosphere (or directly through the star, depending on the geometry of the twist). The current can flow across the field lines in the deep crust because it is able to sustain a significant Ampère force $\mathbf{j} \times \mathbf{B}/c \neq 0$ — this force is balanced by elastic forces of the deformed crust.

The emerging currents are easily maintained during the X-ray outburst accompanying a starquake. A dense, thermalized plasma is then present in the magnetosphere, which easily conducts the current. Plasma remains suspended for some time after the starquake because of the transient thermal afterglow [9]. Eventually the afterglow extinguishes and the radiative flux becomes unable to support the plasma outside the star. The decreasing density then threats the capability of the magnetosphere to conduct the current of the magnetic twist. A minimal "corotation" charge density $\rho_{\rm co} = -\mathbf{\Omega} \cdot \mathbf{B}/2\pi c$ is always maintained [10], but it is far smaller than needed to supply the current $\mathbf{j} \sim (c/4\pi)(B/R_{\rm NS})$. Can the lack of charges stop the flow of electric current? A simple estimate shows that the current cannot stop quickly because of its self-induction. A slow decay of the current generates a sufficient self-induction voltage that helps the magnetosphere to re-generate the plasma that carries the current, so that the twisted force-free configuration persists with a qausi-steady $\mathbf{j} \neq 0$.

The stored energy of non-potential (toroidal) magnetic field associated with the ejected current is subject to gradual dissipation. In our model, this dissipation feeds the observed activity of the corona. The stored energy is concentrated near the star and carried by closed magnetic lines with a maximum extension radius $R_{\max} \sim 2R_{NS}$. Thus, most of the twist energy will be released if these lines untwist, and we focus here on the near magnetosphere $R_{\max} \sim 2R_{NS}$.

Consider a magnetic flux tube with cross section $S \lesssim R_{\rm NS}^2$ and length $L \sim R_{\rm NS}$ which carries a current $I = Sj$. The stored magnetic energy of the current per unit length of the tube is

$$\frac{\mathcal{E}_{twist}}{L} \sim \frac{I^2}{c^2} S. \tag{1}$$

The decay of this energy is associated with an electric field parallel to the magnetic lines $\mathbf{E}_\parallel$: this field can accelerate particles and convert the magnetic energy into plasma energy. Conservation of energy can be expressed as

$$\frac{\partial}{\partial t}\left(\frac{B^2}{8\pi}\right) = -\,\mathbf{E}\cdot\mathbf{j} - \nabla\cdot\left(c\frac{\mathbf{E}\times\mathbf{B}}{4\pi}\right), \tag{2}$$

as follows from Maxwell's equations with $E \ll B$. The first term on the right-hand side is the Joule dissipation caused by $\mathbf{E}_\parallel$. The second term — the divergence of the Poynting flux — vanishes for the induced field $\mathbf{E}_\parallel$.

The decay of the twist is related to the voltage between the footpoints of a magnetic line,

$$\Phi_e = \int_A^C \mathbf{E}\cdot \mathrm{d}\mathbf{l}. \tag{3}$$

Here A and C are the anode and cathode footpoints and d$\mathbf{l}$ is the line element; the integral is taken along the magnetic line outside the star (Fig. 1). The product $\Phi_e I$ approximately represents the dissipation rate $\dot{\mathcal{E}}$ in the tube. (The two quantities are not exactly equal in a time-dependent circuit.) The instantaneous dissipation rate is given by

$$\dot{\mathcal{E}} = \int_A^C \mathbf{E}\cdot\mathbf{I}(l)\mathrm{d}\mathbf{l}, \tag{4}$$



Fig. 1 Schematic picture of a current-carrying magnetic tube anchored to the star's surface. The current is initiated by a starquake that twists one (or both) footpoints of the tube. The current flows along the tube up to the magnetosphere and comes back to the star at the other footpoint. A self-induction voltage is created along the tube between its footpoints, which accompanies the gradual decay of the current. The voltage is generated because the current has a tendency to become charge-starved above the atmospheric layer whose scale-height $h = k_{\rm B}T/gm_p$ is a few cm.

where $\mathbf{I}(l) = S\mathbf{j}$ is the instantaneous current at position l in the tube. The current is fixed at footpoints A and C, $I = I_0$; it may, however, fluctuate between the footpoints.

The voltage Φ_e is entirely maintained by the self-induction that accompanies the gradual untwisting of the magnetic field. Thus, Φ_e reflects the gradual decrease of the magnetic helicity in the tube,

$$\dot{\mathcal{H}} = \frac{\partial}{\partial t}\int_A^C \mathbf{A}\cdot\mathbf{B}\,\mathrm{d}V = -\int_A^C \mathbf{E}\cdot\mathbf{B}\,S(l)\,\mathrm{d}l. \tag{5}$$

(Here $\mathrm{d}V = S(l)\mathrm{d}l$ is the volume element.) It also determines the effective resistivity of the tube: $\mathcal{R} = \Phi_e/I$. The self-induction voltage passes the released magnetic energy to charged particles, and a higher rate of untwisting implies a higher energy $e\Phi_e$ gained per particle. A huge magnetic energy is stored in the twisted tube, and a quick untwisting would lead to extremely high Lorentz factors of the accelerated particles.

There is, however, a bottleneck that prevents a fast decay of the twist: the tube responds to high voltages through the copious production of $e^\pm$ pairs. Runaway pair creation is ignited when electrons are accelerated to a certain Lorentz factor $\gamma_\pm \sim 10^3$ (see below). The created $e^\pm$ plasma does not tolerate large $E_\parallel$ — the plasma is immediately polarized, screening the electric field. This provides a negative feedback that limits the magnitude of Φ_e and buffers the decay of the twist.

A minimum Φ_e is needed for the formation of a corona. Two mechanisms can supply plasma: (1) Ions and electrons can be lifted into the magnetosphere from the stellar surface. This requires a minimum voltage,

$$e\Phi_e \sim g m_i R_{\rm NS} \sim 200\ \ \mathrm{MeV}, \tag{6}$$

corresponding to $E_\parallel$ that is strong enough to lift ions (of mass m_i) from the anode footpoint and electrons from the cathode footpoint. (2) Pairs can be created in the magnetosphere if

$$e\Phi_e \sim \gamma_\pm mc^2 = 0.5\gamma_\pm \ \mathrm{MeV}, \tag{7}$$

which can accelerate particles to the Lorentz factor $\gamma_\pm$ sufficient to ignite $e^\pm$ creation.

If Φ_e is too low and the plasma is not supplied, a flux tube is guaranteed to generate a stronger electric field. The current becomes slightly charge-starved: that is, the net density of free charges becomes smaller than $|\nabla\times\mathbf{B}|/4\pi e$. The ultrastrong magnetic field, whose twist carries an enormous energy compared with the energy of the plasma, does not change and responds to the decrease of $\mathbf{j}$ by generating an electric field: a small reduction of the conduction current $\mathbf{j}$ induces a displacement current $(1/4\pi)\partial\mathbf{E}/\partial t = (c/4\pi)\nabla\times\mathbf{B} - \mathbf{j}$. The longitudinal electric field $E_\parallel$ then quickly grows until it can pull particles away from the stellar surface and ignite pair creation.

The limiting cases $E_\parallel \to 0$ (no decay of the twist) and $E_\parallel \to \infty$ (fast decay) both imply a contradiction. The electric field and the plasma content of the corona must regulate each other toward a self-consistent state, and the gradual decay of the twist proceeds through a delicate balance: $E_\parallel$ must be strong enough to supply plasma and maintain the current in the corona; however, if plasma is oversupplied $E_\parallel$ will be reduced by screening. A cyclic behavior is possible, in which plasma is periodically oversupplied and $E_\parallel$ is screened. Our numerical experiment shows that such a cyclic behavior indeed takes place.

3 Electric circuit: numerical experiment

Three facts facilitate the simulation of the coronal electric circuit:

1. The ultrastrong magnetic field makes the particle dynamics 1-D. The magnetic field lines are not perturbed significantly by the plasma inertia, and they can be thought of as fixed “rails” along which the particles move. The particle motion is confined to the lowest Landau level and is strictly parallel to the field (the lifetime of a particle in an excited Landau state is tiny).

2. The particle motion is collisionless in the magnetosphere. It is governed by two forces only: the component of gravity projected onto the magnetic field and a collective electric field $E_\parallel$ which is determined by the charge density distribution and must be found self-consistently.

3. The star possesses a dense and thin atmospheric layer.[1] Near the base of the atmosphere, the required current $\mathbf{j}_B = (c/4\pi)\nabla\times\mathbf{B}$ is easily conducted, with almost no electric field. Therefore the circuit has simple boundary conditions: $E_\parallel = 0$ and fixed current. The atmosphere is much thicker than the skin depth of the plasma and screens the magnetospheric electric field from the star.

Our goal is to understand the plasma behavior above the screening layer, where the atmospheric density is exponentially reduced and an electric field $E_\parallel$ must develop. The induced electric field $\mathbf{E}$ and conduction current $\mathbf{j}$ satisfy the Maxwell equation,

$$\nabla\times\mathbf{B} = \frac{4\pi}{c}\mathbf{j} + \frac{1}{c}\frac{\partial\mathbf{E}}{\partial t}. \tag{8}$$

Here $\mathbf{j}$ is parallel to the direction of the magnetic field, and the force-free condition requires $\nabla\times\mathbf{B} \parallel \mathbf{B}$. Therefore, in the fixed magnetic configuration where $\nabla\times\mathbf{B}$ does not vary with time, $\partial\mathbf{E}/\partial t$ is parallel to $\mathbf{B}$, i.e. only $E_\parallel$ is created by the self-iduction effect. We consider a simple model problem where no pre-existing perpendicular field $\mathbf{E}_\perp$ exists; more precisely, we require $\nabla\cdot\mathbf{E}_\perp = 0$. Then Gauss' law reads

$$4\pi\rho = \nabla\cdot\mathbf{E}_\parallel = \frac{\mathrm{d}E_\parallel}{\mathrm{d}l}, \tag{9}$$

where l is length measured along the magnetic tube. Then the problem becomes strictly 1-D since $\mathbf{E}_\perp$ has no relation to charge density and falls out from the problem. Note that the approximation of 1D circuit, $j_B(t) = const$, where $j_B \equiv (c/4\pi)|\nabla\times\mathbf{B}|$, excludes the excitation of transverse waves in the magnetosphere, which in reality can exist. These waves are described by the coupled fluctuations δj_B and $\mathbf{E}_\perp$ on scales much smaller than the circuit size L. (Wavelengths $\lambda \sim c/\omega_P$ can be excited by plasma oscillations.) These fluctuations may be expected to have a small effect on the circuit solution unless δj_B becomes comparable to j_B.

With $j_B(t) = const$, the particle motions on different magnetic lines are decoupled. Indeed, the particle dynamics is controlled by electric field $E = E_\parallel$ which is related to the instantaneous charge distribution by Gauss' law (9). Thus, $E(l)$ on a given magnetic line is fully determined by charge density $\rho(l)$ *on the same line*. The plasma and electric field evolve along the line as if the world were 1-D.

If the conduction current j is smaller than j_B, then $\partial E_\parallel/\partial t > 0$ and an electric field appears that tends to increase the current. Alternatively, if $j > j_B$ then an electric field of the opposite sign develops which tends to reduce the current. Thus, j is always regulated toward j_B. This is the standard self-induction effect. The timescale of the regulation, τ, is very short, of the order of the Langmuir oscillation timescale $\omega_P^{-1} \sim 10^{-13}$ s. Local deviations from charge neutrality tend to be erased on the same plasma timescale. The net charge imbalance of the corona must be extremely small — a tiny imbalance would create an electric field that easily pulls out the missing charges from the surface. When the corotation charge density is neglected, the net charge of the corona vanishes.

---

[1] Such a layer initially exists on the surface of a young neutron star. Its maintenance is discussed in [4].



Fig. 2 Set up of numerical experiment. Thin and dense plasma layers are maintained near the cathode and anode by injecting cold particles through the boundaries of the tube (footpoints of a magnetic flux tube). The electric current is kept constant at the boundaries and the system is allowed to evolve in time until a quasi-steady state is reached. Without voltage between anode and cathode, the current cannot flow because gravity g traps the particles near the boundaries. The constant current at the boundaries, however, implies that voltage is immediately generated should the flow of charge stop in the tube. The experiment aims to find the self-induction voltage that keeps the current flowing.



Fig. 3 Circuit without $e^\pm$ production. Upper panel shows the normalized charge density and lower panel shows the electric field in units of E_g defined by $eE_g = m_i g_0$. Solid and dotted curves correspond to two different moments of time, demonstrating that a quasi-steady state has been reached. In this simulation, $m_i = 10 m_e$ and $\zeta = c/\omega_P L \simeq 0.01$. The final state is the relativistic double layer described analytically in [11].

Adopting a characteristic velocity $v \sim c$ of the coronal particles, the Debye length of the plasma $\lambda_{\rm D} = v/\omega_P$ can be taken equal to the plasma skin depth c/ω_P. An important physical parameter of the corona is the ratio of the Debye length to the circuit size $L \sim R_{\rm NS}$,

$$\zeta = \frac{\lambda_{\rm D}}{L} = \frac{c}{L\omega_P} \ll 1. \tag{10}$$

The force-free condition $\mathbf{j}_B \times \mathbf{B}$ together with $\nabla \cdot \mathbf{j}_B = 0$ requires that $j_B(l) \propto B(l)$ along the magnetic line. We can scale out this variation simply dividing all local quantities (charge density, current density, and electric field) by B. This reduces the problem to an equivalent problem where $j_B(l) = const$. Furthermore, only forces along magnetic lines control the plasma dynamics, and the curvature of magnetic lines falls out from the problem. Therefore, we can set up the experiment so that plasma particles move along a straight line connecting anode and cathode (Fig. 2). We designated this line as the z-axis, so that $l = z$.

This one-dimensional system may be simulated numerically. Consider N particles moving along the z-axis ($N \sim 10^6$ in our simulations). The electric field acting on a particle at a given z is given by

$$E(z) = 4\pi \int_0^z \rho \, {\rm d}z, \tag{11}$$

where $E(0) = E(L) = 0$ are the boundary conditions. Given the instantaneous positions z_i of all charges e_i we immediately find $E(z)$. To find a quasi-steady state of such a system we let it relax by following the particle motion in the self-consistent electric field.

Consider first a circuit where $e^\pm$ production is (artificially) forbidden, so that only the cold hydrostatic atmosphere can supply particles to the corona. We find that such a circuit quickly relaxes to a state where it acts as an ultra-relativistic linear accelerator (Fig. 3). In this state, oscillations of electric field (on the plasma timescale ω_P^{-1}) are confined to the thin atmospheric layers. A static accelerating electric field is created above the layers where the atmosphere density is exponentially reduced. The asymmetry of the solution is caused by the difference between the electron and ion masses.

This configuration is a relativistic double layer [11]. It is well described by Carlqvist's solution, which has no gravity in the circuit and assumes zero temperature at the boundaries, so that particles are injected with zero velocity. According to this solution, the potential drop between anode and cathode, in the limit $e\Phi_e \gg m_i c^2$, is

$$e\Phi_e = \left[\left(\frac{m_i}{Z m_e}\right)^{1/2} + 1\right] \frac{L\omega_P}{2}\, m_e c, \qquad \omega_P^2 \equiv \frac{4\pi j e}{m_e c}, \tag{12}$$

where Z is the ion charge number (in our experiment $Z = 1$ was assumed). The established voltage is much larger than is needed to overcome the gravitational barrier Φ_g. It does not even depend on Φ_g as long as Φ_g is large enough to prohibit the thermally-fed regime. Gravity causes the transition to the linear-accelerator state, but the state itself does not depend on Φ_g.

Thus, the first result of numerical experiment is that an ion-electron circuit (without $e^\pm$ creation) in a gravitational field relaxes to the double layer of macroscopic size L and huge voltage Φ_e. The system does not find any state with a lower Φ_e, even though it is allowed to be time-dependent and strong fluctuations persist in the atmospheric layers (reversing the sign of E).

If the linear accelerator were maintained in the twisted magnetosphere, the twist would be immediately killed off: the huge voltage implies a large untwisting rate (§ 2). The electron Lorentz factor developed in the linear accelerator is (taking the real $m_i/m_e = 1836$ and $\zeta \sim 3 \times 10^{-9}$),

$$\gamma_e = \frac{e\Phi_e}{m_e c^2} \simeq \frac{20}{\zeta} \sim 6 \times 10^9. \qquad (13)$$

However, new processes will become important before the particles could acquire such high energies: production of $e^\pm$ pairs will take place. Therefore, the linear-accelerator solution cannot describe a real magnetosphere. We conclude that pair creation is a key ingredient of the circuit that will regulate the voltage to a smaller value.



Fig. 4 Spacetime diagrams illustrating the critical character of the $e^\pm$ breakdown and formation of avalanches. The seed electron is placed at cathode and is accelerated to $\gamma_{\rm res}$ after passing the distance $a_{\rm res} = (\gamma_{\rm res} - 1)m_e c^2/eE$. The figure shows the worldlines of created particles. (a) $a_{\rm res}/L = 0.35$ (supercritical case). (b) $a_{\rm res}/L = 0.6$ (critical case).

4 Pair discharge and self-organized criticality

In the magnetospheres of canonical radio pulsars with $B \sim 10^{12}$ G, $e^\pm$ pairs are created when seed electrons are accelerated to large Lorentz factors $\gamma_e \sim 10^7$. Such electrons emit curvature gamma rays that can convert to $e^\pm$ off the magnetic field. In stronger fields, another channel of $e^\pm$ creation appears. It is also two-step: an accelerated particle resonantly upscatters a thermal X-ray photon, which subsequently converts to a pair [12, 4]. The resonant scattering requires the electron to have a Lorentz factor,

$$\gamma_{\rm res} = \frac{B/B_{\rm QED}}{1 - \cos\theta_{kB}} \, \frac{m_e c^2}{\hbar\omega_X} \approx 10^3 \, B_{15} \, \frac{10 \text{ keV}}{\hbar\omega_X}, \qquad (14)$$

where $\hbar\omega_X$ and θ_{kB} are the energy and pitch angle of the target photon. The distance over which an accelerated electron creates a pair is approximately equal to the free path to scattering $l_{\rm res}$. It depends only on B and the spectrum of target photons; $l_{\rm res} \ll R_{\rm NS}$ for the relevant parameter $B_{15}\hbar\omega_X < 100$ keV [4].

When the voltage in the circuit becomes high enough to accelerate electrons to $\gamma_{\rm res}$, an $e^\pm$ breakdown develops in the magnetic flux tube. An illustrative toy model of breakdown is shown in a spacetime diagram in Figure 4. Each pair-creation event gives two new particles of opposite charge, which initially move in the same direction. One of them is accelerated by the electric field and lost after a time $< L/c$, while the other is decelerated and can reverse direction before reaching the boundary. This reversal of particles in the tube and repeated pair creation allows the $e^\pm$ plasma to be continually replenished. In the super-critical regime (left panel in Fig. 4) more than one reversing particle is created per passage time L/c and an avalanche develops exponentially on a timescale $\sim a_{\rm res}/c$. In the near-critical regime shown in the right panel, just one reversing particle is created per passage time L/c. This critical state is unstable: sooner or later the avalanche will be extinguished.

This toy model shows an essential property of the $e^\pm$ breakdown: it is a critical stochastic phenomenon. Above a critical voltage pair creation proceeds in a runway manner, and the current and the dissipation rate would run away if the voltage were fixed. Below the critical voltage pair creation does not ignite. The criticality parameter is $L/a_{\rm res} = e\Phi_e/(\gamma_{\rm res} - 1)m_e c^2$. The tube with enforced current at the boundaries must self-organize to create pairs in the near-critical regime $e\Phi_e \sim \gamma_{\rm res} m_e c^2$ and maintain the current.

The discharging tube is similar to other phenomena that show self-organized criticality, e.g., a pile of sand on a table [13]. If sand is steadily added, a quasi-steady state is established with a characteristic mean slope of the pile. The sand is lost (falls from the table) intermittently, through avalanches — a sort of "sand discharge." In our case, charges of the opposite signs are added steadily instead of sand (fixed j at the boundaries), and voltage $\Phi_e = EL$ plays the role of the mean slope of a pile. The behavior of the discharging system is expected to be time-dependent, with stochastic avalanches.

We implemented the process of pair creation in our numerical experiment. Regardless the details of this process (values of $\gamma_{\rm res}$ and $l_{\rm res}$) the circuit relaxed to the state of self-organized criticality after a few $t_{\rm dyn} = L/c$. This state is time-dependent on short timescales, but it has a well-defined steady voltage when averaged over a few $t_{\rm dyn}$ (Fig. 5),

$$e\bar{\Phi}_e \sim \gamma_{\rm res} m_e c^2. \quad (15)$$

We found that this relation applies even to circuits with $\gamma_{\rm res} m_e c^2 \gg m_i c^2$, where lifting of the ions is energetically preferable to pair creation. We conclude that *the voltage along the magnetic tube is self-regulated to a value just enough to maintain pair production and feed the current with $e^\pm$ pairs.* In all cases, the pair creation rate $2\dot{N}_+ \sim j_B/e$ is maintained.

The solution for the plasma dynamics in the corona is strongly non-linear and time-dependent. It is essentially global in the sense that the plasma behavior near one footpoint of a magnetospheric field line is coupled to the behavior near the other footpoint. Remarkably, this complicated global behavior may be described as essentially one-dimensional electric circuit that is subject to simple boundary conditions $E \approx 0$ on the surface of the star and can be studied using a direct numerical experiment.

The current is carried largely by $e^\pm$ everywhere in the tube. The ion fraction in the current depends on the ratio $e\Phi_e/m_i c^2 \simeq \gamma_{\rm res} m_e/m_i$ (Fig. 6). If this ratio is small, pairs are easily produced with a small electric field, too small to lift the ions, and the ion current is suppressed. In the opposite case, ions carry about 1/2 of the current. By coincidence, $\gamma_{\rm res} \lesssim m_i/m_e$ is typical for magnetars, which implies that ions carry $\lesssim 10\%$ of the current.

5 Implications

The basic finding of this work is that an $e^\pm$ corona must be maintained around a magnetar. It exists in a state of self-organized criticality, near the threshold for $e^\pm$ breakdown. The stochastic $e^\pm$ discharge continually replenishes the coronal plasma, which is lost on the dynamic time $t_{\rm dyn} \sim 10^{-4}$ s. The established voltage along a twisted magnetic tube is marginally sufficient to accelerate an electron (or positron) to the energy where collisions with ambient X-ray photons spawn new pairs. This physical condition determines the rate of energy dissipation in the corona. The generation of this voltage is precisely the self-induction effect of the gradually decaying magnetic twist, and the energy release in the corona is fed by the magnetic energy of the twisted field.

The rate of energy dissipation in the twisted magnetosphere is given by $L_{\rm diss} = I\Phi_e$ where I is the net current through the corona. The current may be estimated as $I \sim j_B a^2 \sim \frac{c}{4\pi}\Delta\phi\,(B/R_{\rm NS})\,a^2$, where a is the size of a twisted region on the stellar surface and $\Delta\phi = |\nabla\times\mathbf{B}|(B/R_{\rm NS})^{-1} \lesssim 1$ characterizes the strength of the twist. The calculated $e\Phi_e \sim 1$ GeV implies

$$L_{\rm diss} = I\Phi_e \sim 10^{37}\,\Delta\phi\,B_{15}\left(\frac{a}{R_{\rm NS}}\right)^2\left(\frac{e\Phi_e}{\rm GeV}\right)\frac{\rm erg}{\rm s}. \quad (16)$$

Observed luminosity $L_{\rm diss} \sim 10^{36}$ erg s^{-1} is consistent with a partially twisted magnetosphere, $a \sim 0.3R_{\rm NS}$, or a global moderate twist with $\Delta\phi\,(e\Phi_e/{\rm GeV}) \sim 0.1$.

Once created, a magnetospheric twist has a relatively long but finite lifetime. The energy stored in it is $\mathcal{E}_{\rm twist} \sim (I^2/c^2)R_{\rm NS}$. This energy dissipates in a time

$$t_{\rm decay} = \frac{\mathcal{E}_{\rm twist}}{L_{\rm diss}} \approx 3\left(\frac{L_{\rm diss}}{10^{36}\ {\rm erg\ s^{-1}}}\right)\left(\frac{e\Phi_e}{\rm GeV}\right)^{-2}\ {\rm yr}. \quad (17)$$

Note that a stronger twist (brighter corona) lives longer. If the time between large-scale starquakes is longer than $t_{\rm decay}$, the magnetar should be seen to enter a quiescent state. Such behavior has been observed in AXP J1810-197: an outburst was followed by the gradual decay on a year timescale [14].



Fig. 5 Time history of one numerical experiment. The circuit parameters are: $m_i = 10m_e$, $\gamma_{\rm res} = m_i/m_e$, $\zeta = 0.01$. After a few dynamical times the voltage Φ_e stops growing and the circuit enters an oscillatory regime. During each oscillation, an increased voltage triggers $e^\pm$ discharge, then $\dot{N}_+$ drops, and Φ_e begins to grow again. The quasi-steady oscillating state (self-organized criticality) is established. The horizontal lines show the voltage $e\Phi_e = \gamma_{\rm res} m_e c^2$ and the minimum pair production rate $2\dot{N}_+ = j_B/e$ that can feed the required current j_B.



Fig. 6 Fraction of the electric current carried by ions vs. voltage. In the left panel voltage is taken in units of $m_i c^2$, and in the right panel — in units of $\gamma_{\rm res} m_e c^2$. Different colors and symbols show circuits with $m_i/m_e = 10, 30$ and various values of $\gamma_{\rm res}$ ranging from $m_i/5m_e$ to $5m_i/m_e$ (see [4]).

The voltage Φ_e implies a certain effective resistivity of the corona, $\mathcal{R} = \Phi_e/I$. This resistivity leads to spreading of the electric current *across* the magnetic lines, which is described by the induction equation $\partial \mathbf{B}/\partial t = -c\nabla \times \mathbf{E}$. The timescale of twist spreading to the light cylinder is comparable to $t_{\rm decay}$. Therefore, the impact of a starquake on spindown may appear with a delay of $\sim$ years.

Even a small ion current implies a large mass transfer through the magnetosphere over the $\sim 10^4$-yr active lifetime of a magnetar. It requires a significant excavation of the crust at the anode footpoints of twisted magnetic tubes. The bombarding relativistic electrons from the corona cause this excavation. They spall heavy ions in the uppermost crust, re-generate the light-element atmosphere on the surface, and regulate its column density to a value $\sim 100 m_p/\sigma_{\rm T}$ [4].

The radiative output from the corona likely peaks in its inner region because the coronal current is concentrated on closed field lines with a maximum extension $R_{\rm max} \lesssim 2R_{\rm NS}$ (§ 2). Emission from the inner corona must be suppressed above ~ 1 MeV, regardless the mechanism of emission, because photons with energy $\gtrsim 1$ MeV cannot escape the ultra-strong magnetic field. The observed high-energy radiation extends above 100 keV [2, 15, 16, 17, 18]. There is an indication for a cutoff between 200 keV and 1 MeV from COMPTEL upper limits for AXP 4U 0142+61.

The possible mechanisms of emission are strongly constrained. The corona has a low density $n \sim n_c = j_B/ec \lesssim B(4\pi\, e R_{\rm NS})^{-1} \sim 10^{17} B_{15}$ cm^{-3} and particle collisions are rare, so two-body radiative processes are negligible. The upscattering of keV photons streaming from the surface does not explain the observed 100-keV emission [4]. A possible source of the 100 keV X-rays is the transition layer between the corona and the thermal photosphere [19, 4]. It is heated by the coronal beam and cooled by heat conduction toward the surface, and its temperature is regulated to $\sim 100(\ell/\ell_{\rm Coul})^{-2/5}$ keV, where $\ell/\ell_{\rm Coul} < 1$ parameterizes the suppression of thermal conductivity by plasma turbulence. The emitted X-ray spectrum may be approximated as single-temperature optically thin bremsstrahlung [19]. Its photon index below the exponential cutoff is close to -1, in agreement with spectra observed with INTEGRAL and RXTE.

The observed pulsed fraction increases toward the high-energy end of the spectrum and approaches 100% at 100 keV (Kuiper et al. 2004). If the hard X-rays are produced by one or two twisted spots on the star surface, the large pulsed fraction implies that the spots almost disappear during some phase of rotation. Then they should not be too far from each other. They cannot be, e.g., antipodal because a large fraction of the star surface is visible to observer due to the gravitational bending of light: $S_{\rm vis}/4\pi R_{\rm NS}^2 = (2 - 4GM/c^2 R_{\rm NS})^{-1} \approx 3/4$ [20].

The transition layer at a footprint of a coronal flux tube may reach high enough temperature for thermal $e^\pm$ production [4]. The created pairs then elevate above the surface and can conduct the magnetospheric current. This opens a possibility for the corona to be fed by surface pair creation instead of the $e^\pm$ discharge.

The observed infrared (K-band) and optical luminosities of magnetars are $\sim 10^{32}$ erg/s [21], which is far above the Rayleigh-Jeans tail of the surface blackbody radiation. The inferred brightness temperature is $\sim 10^{13}$ K if the radiation is emitted from the surface of the star. Four principle emission mechanisms may then be considered: coherent plasma emission, synchrotron emission by electrons, cyclotron emission by ions, and curvature radiation by $e^\pm$ bunches in the corona. The last two are most plausible (see [4] for detailed discussion). In particular, curvature radiation in the inner magnetosphere may extend into the optical-IR range. Charges moving on field lines with a curvature radius $R_C \lesssim R_{\rm NS}$ will emit radiation with frequency $\nu_C \sim \gamma_e^3 c/2\pi R_C$, which is in the K-band ($\nu = 10^{14}$ Hz) or optical when $\gamma_e m_e c^2 \gtrsim$ GeV. The radio luminosity of a normal pulsar can approach $\sim 10^{-2}$ of its spindown power, and it is not implausible that the observed fraction $\sim 10^{-4}$ of the power dissipated in a magnetar corona is radiated in optical-IR photons by the same coherent mechanism.

The corona luminosity is simply proportional to the net current flowing through the magnetosphere, and is determined by the competition between occasional twisting (due to starquakes) and gradual dissipative untwisting of the magnetosphere. During periods of high activity, when starquakes occur frequently, the magnetospheric twist may grow to the point of a global instability: when a critical $\Delta\phi \gtrsim 1$ is achieved, the magnetosphere suddenly relaxes to a smaller twist angle. A huge release of energy must then occur, producing a giant flare; a model of how this can happen is discussed in [22].

A possible way to probe the twist evolution is to measure the history of spindown rate $\dot{P}$ of the magnetar. The existing data confirm the theoretical expectations: the spin-down of SGRs was observed to accelerate months to years following periods of activity [23,24]. A similar effect was also observed following flux enhancement in AXPs [25]. This "hysteresis" behavior of the spindown rate may represent the delay with which the twist is spreading to the outer magnetosphere.

Since the density of the plasma corona is proportional to the current that flows through it, changes in the twist also lead to changes in the coronal opacity. The density of the plasma corona is close to its minimum $n_c = j/ec$, and it is largely made of $e^{\pm}$ pairs rather than electron-ion plasma. The opacity should increase following bursts of activity and affect the X-ray pulse profile and spectrum. Such changes, which persist for months to years, have been observed following X-ray outbursts from two SGRs [26].

Acknowledgements This work was supported by NASA grant NNG-06-G107G.

References

1. Woods, P. M., & Thompson, C. 2006, in "Compact Stellar X-ray Sources," eds. W. H. G. Lewin and M. van der Klis, Cambridge University Press, astro-ph/0406133
2. Kuiper, L., Hermsen, W., den Hartog, P. R., Collmar, W. 2006, ApJ, 645, 556
3. Thompson, C., Lyutikov, M., & Kulkarni, S. R. 2002, ApJ, 574, 332
4. Beloborodov, A. M., & Thompson, C. 2006, ApJ, submitted (astro-ph/0602417)
5. Ruderman, M. A., & Sutherland, P. G. 1975, ApJ, 196, 51
6. Arons, J., & Scharlemann, E. T. 1979, ApJ, 231, 854
7. Thompson, C., & Duncan, R. C. 2001, ApJ, 561, 980
8. Thompson, C., & Duncan, R. C. 1995, MNRAS, 275, 255
9. Ibrahim, A. I., et al. 2001, ApJ, 558, 237
10. Goldreich, P., & Julian, W. H. 1969, ApJ, 157, 869
11. Carlqvist, P. 1982, Ap&SS, 87, 21
12. Hibschman, J. A., & Arons, J. 2001, ApJ, 554, 624
13. Bak, P., Tang, C., & Wiesenfeld, K. 1987, Phys. Rev. Lett., 59, 381
14. Gotthelf, E. V., & Halpern, J. P. 2005, ApJ, 632, 1075
15. Kuiper, L., Hermsen, W., & Mendez, M. 2004, ApJ, 613, 1173
16. Mereghetti, S., Götz, D., Mirabel, I. F., & Hurley, K. 2005, A&A, 433, L9
17. Molkov, S., et al. 2005, A&A, 433, L13
18. den Hartog, P. R., et al. 2006, A&A, in press (astro-ph/0601644)
19. Thompson, C., & Beloborodov, A. M. 2005, ApJ, 634, 565
20. Beloborodov, A. M. 2002, ApJ, 566, L85
21. Hulleman, F., van Kerkwijk, M. H., & Kulkarni, S. R. 2004, A&A, 416, 1037
22. Lyutikov, M. 2006, MNRAS, 367, 1594
23. Kouveliotou, C., et al. 1999, ApJ, 510, L115
24. Woods, P. M., et al. 2002, ApJ, 576, 381
25. Gavriil, F. P., & Kaspi, V. M. 2004, ApJ, 609, L67
26. Woods, P. M., et al. 2001, ApJ, 552, 748